

CABINET:

Nighttime Severe Cold & Flu



Active Ingredients (in each softgel)

Purpose

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • cough to help you sleep • minor aches & pains • headache • fever • sore throat • runny nose & sneezing • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product. • to make a child sleep

Ask a doctor or pharmacist before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin

When using this product • do not use more than directed

• excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

• take only as directed – see **Overdose warning** • do not exceed 4 doses per 24 hours • adults & children 12 years & over; 2 soft gels with water every 4 hours • children 4 to under 12 years; ask a doctor • children under 4 years; do not use

Other Information

• store at room temperature

Inactive Ingredients

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol 400, povidone K30, propylene glycol, purified water, shellac, sorbitol sorbitan, sodium hydroxide, titanium dioxide