

CABINET:

Stool Softener



Active Ingredients (in each softgel)

Purpose

Docusate sodium 100 mg.....Stool softener

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 12-72 hours

Warnings

Do not use if you are presently taking mineral oil, unless told to do so by a doctor

Ask a doctor before use if you have • stomach pain • nausea • vomiting • noticed a sudden change in bowel habits that lasts over 2 weeks

Stop use and ask a doctor if • you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition • you need to use a stool softener laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- take only by mouth. Doses may be taken as a single daily dose or in divided doses. • adults & children 12 years of age & over; take 1-3 soft gels daily • children 2 to under 12 years of age; take 1 soft gel daily • children under 2 years of age; ask a doctor

Other Information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) • Keep tightly closed

Inactive Ingredients

FD&C Red #40, FD&C Yellow #6, gelatin, glycerin, polyethylene glycol-400, povidone, propylene glycol, purified water, sorbitol solution, titanium dioxide