

CABINET:

Laxative



Active Ingredients (in each tablet)	Purpose
Bisacodyl (USP) 5 mg.....	Stimulant Laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6-12 hours

Warnings

Do not use if you cannot swallow without chewing

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

When using this product • do not chew or crush tablet(s)

- it may cause stomach discomfort, faintness and cramps
- do not use within 1 hour after taking an antacid or milk

Stop use and ask a doctor if • you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition • you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- take with a glass of water
- adults & children 12 years of age & over: 1 to 3 tablets in a single daily dose
- children 6 to under 12 years: 1 tablet in a single daily dose
- children under 6 years of age: ask a doctor

Other Information

Each tablet contains: magnesium 5 mg • store at 20-25°C (68-77°F) • protect from excessive humidity

Inactive Ingredients

Acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C Red #27 aluminum lake, D&C Yellow #10 aluminum lake, FD&C Blue #2 aluminum lake, FD&C Yellow 6, gelatin, iron oxide, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, poly ethylene glycol, polyvinyl acetate phtalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, tale, titanium dioxide laurylsulphate, stearic acid