

CABINET:

Pain Reliever



Active Ingredients (in each tablet)

Purpose

Aspirin 81 mg (NSAID)*Pain reliever
*nonsteroidal anti-inflammatory drug

Uses

• for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches, fever or other symptoms needing immediate relief.** • ask your doctor about other uses for Aspirin 81 mg

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or non prescription NSAIDs [aspirin, ibuprofen, naproxen, or others] • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for: • gout • diabetes • arthritis

Stop use and ask a doctor if • an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • pain gets worse or lasts more than 10 days • redness or swelling is present • new symptoms occur • ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose,** get medical help or contact a Poison Control Center right away.(1-800-222-1222)

Directions

• drink a full glass of water with each dose • adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor • children under 12 years: consult a doctor

Other Information

• **each tablet contains:** sodium 0.36 mg • store at room temperature

Inactive Ingredients

Acrycoat, colloidal silicon dioxide, croscarmellose sodium, D&C Yellow No. 10, lactose, microcrystalline cellulose, polysorbate 80, sodium hydroxide, talc, titanium dioxide, triethyl citrate, yellow oxide of iron