

CABINET:

Extra Strength Headache Relief



Active Ingredients (in each tablet)

Purpose

Acetaminophen 250 mg.....	Pain reliever
Aspirin 250 mg (NSAID)*.....	Pain reliever
Caffeine 65 mg.....	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
- headache
- a cold
- arthritis
- muscular aches
- sinusitis
- toothache
- premenstrual & menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away. Aspirin may cause a severe allergic reaction which may include: hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 8 tablets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks everyday while using this product • take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if • you have liver disease • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis or kidney disease • you are taking a diuretic • you have asthma

Ask a doctor or pharmacist before use if you are taking • any other drug containing an NSAID (prescription or non-prescription) • a prescription drug for diabetes, gout or arthritis • any other drug, or are under a doctor's care for any serious condition • a blood thinning (anticoagulant) or steroid drug

Stop use and ask a doctor if • an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • ringing in the ears or loss of hearing occurs • painful area is red or swollen • pain gets worse or lasts for more than 10 days • fever gets worse or lasts for more than 3 days • any new symptoms appear. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not use more than directed (see Overdose warning)
- drink a full glass of water with each dose • adults and children 12 years and over: take 2 tablets every 6 hours: not more than 8 tablets in 24 hours • children under 12 years: ask a doctor

Other Information

- store below 25°C (77°F) • Tamper Evident Feature: Do not use • store at controlled room temperature 20°-25° C (68°-77° F) • read all product information before using

Inactive Ingredients

Colloidal silicon dioxide, hyper mellose, micro crystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, talc