

CABINET:

Pain Reliever & Nighttime Sleep Aid



Active Ingredients (in each tablet)	Purpose
Acetaminophen 500 mg.....	Pain reliever
Diphenhydramine HCl 25 mg.....	Nighttime Sleep aid

Uses

- temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use • with other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • with any other product containing diphenhydramine, even one used on skin • in children under 12 years of age • if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have • liver disease • a breathing problem such as emphysema or chronic bronchitis • trouble urinating due to an enlarged prostate gland • glaucoma

Ask a doctor or pharmacist before use if you are • taking the blood thinning drug warfarin • taking sedatives or tranquilizers

When using this product • drowsiness will occur • avoid alcoholic drinks • do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if • sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. • pain gets worse or lasts for more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than recommended** (see **Overdose warning**)
- adults & children 12 years & over: • take 2 tablets at bedtime: • do not take more than: 2 tablets of this product in 24 hours
- children under 12 years: do not use this adult product in children under 12 years of age: this will provide more than the recommended dose (overdose) and may cause liver damage

Other Information

- **each tablet contains:** magnesium 0.05 mg
- store between 20-25°C (68-77°F)

Inactive Ingredients

FD&C Blue No. 1, FD&C Blue No. 2, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone K-30, pregelatinized starch, purified water, stearic acid, titanium dioxide